

# AP Calculus BC

## Summer Work

### Instructions

1. Send Dr. Durand an email with your summer contact information – (phone, email) – [durand@lisaacademy.org](mailto:durand@lisaacademy.org)
2. Work all problems, showing all work.
3. If you have questions, email Dr. Durand immediately so that he can help you.

**IMPORTANT REMINDER:** You **MUST** complete your summer assignment in order to be permitted to take the AP Calculus BC course in the Fall.

**2007 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)**

*calculator permitted*

3. The wind chill is the temperature, in degrees Fahrenheit ( $^{\circ}\text{F}$ ), a human feels based on the air temperature, in degrees Fahrenheit, and the wind velocity  $v$ , in miles per hour (mph). If the air temperature is  $32^{\circ}\text{F}$ , then the wind chill is given by  $W(v) = 55.6 - 22.1v^{0.16}$  and is valid for  $5 \leq v \leq 60$ .
- (a) Find  $W'(20)$ . Using correct units, explain the meaning of  $W'(20)$  in terms of the wind chill.
- (b) Find the average rate of change of  $W$  over the interval  $5 \leq v \leq 60$ . Find the value of  $v$  at which the instantaneous rate of change of  $W$  is equal to the average rate of change of  $W$  over the interval  $5 \leq v \leq 60$ .
- (c) Over the time interval  $0 \leq t \leq 4$  hours, the air temperature is a constant  $32^{\circ}\text{F}$ . At time  $t = 0$ , the wind velocity is  $v = 20$  mph. If the wind velocity increases at a constant rate of 5 mph per hour, what is the rate of change of the wind chill with respect to time at  $t = 3$  hours? Indicate units of measure.
- 

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**END OF PART A OF SECTION II**


2007 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

CALCULUS AB  
SECTION II, Part B

Time—45 minutes

Number of problems—3


No calculator is allowed for these problems.



4. Let  $f$  be a function defined on the closed interval  $-5 \leq x \leq 5$  with  $f(1) = 3$ . The graph of  $f'$ , the derivative of  $f$ , consists of two semicircles and two line segments, as shown above.

- For  $-5 < x < 5$ , find all values  $x$  at which  $f$  has a relative maximum. Justify your answer.
- For  $-5 < x < 5$ , find all values  $x$  at which the graph of  $f$  has a point of inflection. Justify your answer.
- Find all intervals on which the graph of  $f$  is concave up and also has positive slope. Explain your reasoning.
- Find the absolute minimum value of  $f(x)$  over the closed interval  $-5 \leq x \leq 5$ . Explain your reasoning.

WRITE ALL WORK IN THE EXAM BOOKLET.

No  
calculator

5. Let  $g$  be a continuous function with  $g(2) = 5$ . The graph of the piecewise-linear function  $g'$ , the derivative of  $g$ , is shown above for  $-3 \leq x \leq 7$ .
- Find the  $x$ -coordinate of all points of inflection of the graph of  $y = g(x)$  for  $-3 < x < 7$ . Justify your answer.
  - Find the absolute maximum value of  $g$  on the interval  $-3 \leq x \leq 7$ . Justify your answer.
  - Find the average rate of change of  $g(x)$  on the interval  $-3 \leq x \leq 7$ .
  - Find the average rate of change of  $g'(x)$  on the interval  $-3 \leq x \leq 7$ . Does the Mean Value Theorem applied on the interval  $-3 \leq x \leq 7$  guarantee a value of  $c$ , for  $-3 < c < 7$ , such that  $g''(c)$  is equal to this average rate of change? Why or why not?

---

**WRITE ALL WORK IN THE EXAM BOOKLET.**

2008 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

CALCULUS AB  
SECTION II, Part A

Time—45 minutes

Number of problems—3

A graphing calculator is required for some problems or parts of problems.

---

1. Let  $R$  be the region in the first quadrant bounded by the graphs of  $y = \sqrt{x}$  and  $y = \frac{x}{3}$ .

- (a) Find the area of  $R$ .
  - (b) Find the volume of the solid generated when  $R$  is rotated about the vertical line  $x = -1$ .
  - (c) The region  $R$  is the base of a solid. For this solid, the cross sections perpendicular to the  $y$ -axis are squares. Find the volume of this solid.
- 

WRITE ALL WORK IN THE EXAM BOOKLET.

Calculator permitted

2008 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

Distance from the river's edge (feet)	0	8	14	22	24
Depth of the water (feet)	0	7	8	2	0

3. A scientist measures the depth of the Doe River at Picnic Point. The river is 24 feet wide at this location. The measurements are taken in a straight line perpendicular to the edge of the river. The data are shown in the table above. The velocity of the water at Picnic Point, in feet per minute, is modeled by  $v(t) = 16 + 2\sin(\sqrt{t+10})$  for  $0 \leq t \leq 120$  minutes.
- (a) Use a trapezoidal sum with the four subintervals indicated by the data in the table to approximate the area of the cross section of the river at Picnic Point, in square feet. Show the computations that lead to your answer.
- (b) The volumetric flow at a location along the river is the product of the cross-sectional area and the velocity of the water at that location. Use your approximation from part (a) to estimate the average value of the volumetric flow at Picnic Point, in cubic feet per minute, from  $t = 0$  to  $t = 120$  minutes.
- (c) The scientist proposes the function  $f$ , given by  $f(x) = 8\sin\left(\frac{\pi x}{24}\right)$ , as a model for the depth of the water, in feet, at Picnic Point  $x$  feet from the river's edge. Find the area of the cross section of the river at Picnic Point based on this model.
- (d) Recall that the volumetric flow is the product of the cross-sectional area and the velocity of the water at a location. To prevent flooding, water must be diverted if the average value of the volumetric flow at Picnic Point exceeds 2100 cubic feet per minute for a 20-minute period. Using your answer from part (c), find the average value of the volumetric flow during the time interval  $40 \leq t \leq 60$  minutes. Does this value indicate that the water must be diverted?

---

WRITE ALL WORK IN THE EXAM BOOKLET.

END OF PART A OF SECTION II

2009 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

*Calculator permitted*


2. A storm washed away sand from a beach, causing the edge of the water to get closer to a nearby road. The rate at which the distance between the road and the edge of the water was changing during the storm is modeled by  $f(t) = \sqrt{t} + \cos t - 3$  meters per hour,  $t$  hours after the storm began. The edge of the water was 35 meters from the road when the storm began, and the storm lasted 5 hours. The derivative of  $f(t)$  is  $f'(t) = \frac{1}{2\sqrt{t}} - \sin t$ .
- (a) What was the distance between the road and the edge of the water at the end of the storm?
  - (b) Using correct units, interpret the value  $f'(4) = 1.007$  in terms of the distance between the road and the edge of the water.
  - (c) At what time during the 5 hours of the storm was the distance between the road and the edge of the water decreasing most rapidly? Justify your answer.
  - (d) After the storm, a machine pumped sand back onto the beach so that the distance between the road and the edge of the water was growing at a rate of  $g(p)$  meters per day, where  $p$  is the number of days since pumping began. Write an equation involving an integral expression whose solution would give the number of days that sand must be pumped to restore the original distance between the road and the edge of the water.

---

WRITE ALL WORK IN THE EXAM BOOKLET.

2009 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

*Calculator permitted*



Graph of  $f$

3. A continuous function  $f$  is defined on the closed interval  $-4 \leq x \leq 6$ . The graph of  $f$  consists of a line segment and a curve that is tangent to the  $x$ -axis at  $x = 3$ , as shown in the figure above. On the interval  $0 < x < 6$ , the function  $f$  is twice differentiable, with  $f''(x) > 0$ .
- Is  $f$  differentiable at  $x = 0$ ? Use the definition of the derivative with one-sided limits to justify your answer.
  - For how many values of  $a$ ,  $-4 \leq a < 6$ , is the average rate of change of  $f$  on the interval  $[a, 6]$  equal to 0? Give a reason for your answer.
  - Is there a value of  $a$ ,  $-4 \leq a < 6$ , for which the Mean Value Theorem, applied to the interval  $[a, 6]$ , guarantees a value  $c$ ,  $a < c < 6$ , at which  $f'(c) = \frac{1}{3}$ ? Justify your answer.
  - The function  $g$  is defined by  $g(x) = \int_0^x f(t) dt$  for  $-4 \leq x \leq 6$ . On what intervals contained in  $[-4, 6]$  is the graph of  $g$  concave up? Explain your reasoning.

**WRITE ALL WORK IN THE EXAM BOOKLET.**

**END OF PART A OF SECTION II**



No calculator

$t$ (seconds)	0	8	20	25	32	40
$v(t)$ (meters per second)	3	5	-10	-8	-4	7

6. The velocity of a particle moving along the  $x$ -axis is modeled by a differentiable function  $v$ , where the position  $x$  is measured in meters, and time  $t$  is measured in seconds. Selected values of  $v(t)$  are given in the table above. The particle is at position  $x = 7$  meters when  $t = 0$  seconds.
- Estimate the acceleration of the particle at  $t = 36$  seconds. Show the computations that lead to your answer. Indicate units of measure.
  - Using correct units, explain the meaning of  $\int_{20}^{40} v(t) dt$  in the context of this problem. Use a trapezoidal sum with the three subintervals indicated by the data in the table to approximate  $\int_{20}^{40} v(t) dt$ .
  - For  $0 \leq t \leq 40$ , must the particle change direction in any of the subintervals indicated by the data in the table? If so, identify the subintervals and explain your reasoning. If not, explain why not.
  - Suppose that the acceleration of the particle is positive for  $0 < t < 8$  seconds. Explain why the position of the particle at  $t = 8$  seconds must be greater than  $x = 30$  meters.


---

WRITE ALL WORK IN THE EXAM BOOKLET.

END OF EXAM

**CALCULUS AB**  
**SECTION II, Part A**  
 Time—45 minutes  
 Number of problems—3

A graphing calculator is required for some problems or parts of problems.



- 1 In the figure above,  $R$  is the shaded region in the first quadrant bounded by the graph of  $y = 4\ln(3 - x)$ , the horizontal line  $y = 6$ , and the vertical line  $x = 2$ .
- Find the area of  $R$ .
  - Find the volume of the solid generated when  $R$  is revolved about the horizontal line  $y = 8$ .
  - The region  $R$  is the base of a solid. For this solid, each cross section perpendicular to the  $x$ -axis is a square. Find the volume of the solid.

---


**WRITE ALL WORK IN THE EXAM BOOKLET.**

2010 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

- ~~X~~ The function  $g$  is defined for  $x > 0$  with  $g(1) = 2$ ,  $g'(x) = \sin\left(x + \frac{1}{x}\right)$ , and  $g''(x) = \left(1 - \frac{1}{x^2}\right)\cos\left(x + \frac{1}{x}\right)$ .
- Find all values of  $x$  in the interval  $0.12 \leq x \leq 1$  at which the graph of  $g$  has a horizontal tangent line.
  - On what subintervals of  $(0.12, 1)$ , if any, is the graph of  $g$  concave down? Justify your answer.
  - Write an equation for the line tangent to the graph of  $g$  at  $x = 0.3$ .
  - Does the line tangent to the graph of  $g$  at  $x = 0.3$  lie above or below the graph of  $g$  for  $0.3 < x < 1$ ? Why?

Calculator permitted

$t$	0	2	4	6	8	10	12
$P(t)$	0	46	53	57	60	62	63



3. The figure above shows an aboveground swimming pool in the shape of a cylinder with a radius of 12 feet and a height of 4 feet. The pool contains 1000 cubic feet of water at time  $t = 0$ . During the time interval  $0 \leq t \leq 12$  hours, water is pumped into the pool at the rate  $P(t)$  cubic feet per hour. The table above gives values of  $P(t)$  for selected values of  $t$ . During the same time interval, water is leaking from the pool at the rate  $R(t)$  cubic feet per hour, where  $R(t) = 25e^{-0.05t}$ . (Note: The volume  $V$  of a cylinder with radius  $r$  and height  $h$  is given by  $V = \pi r^2 h$ .)
- Use a midpoint Riemann sum with three subintervals of equal length to approximate the total amount of water that was pumped into the pool during the time interval  $0 \leq t \leq 12$  hours. Show the computations that lead to your answer.
  - Calculate the total amount of water that leaked out of the pool during the time interval  $0 \leq t \leq 12$  hours.
  - Use the results from parts (a) and (b) to approximate the volume of water in the pool at time  $t = 12$  hours. Round your answer to the nearest cubic foot.
  - Find the rate at which the volume of water in the pool is increasing at time  $t = 8$  hours. How fast is the water level in the pool rising at  $t = 8$  hours? Indicate units of measure in both answers.

WRITE ALL WORK IN THE EXAM BOOKLET.


END OF PART A OF SECTION II

CALCULUS AB  
SECTION II, Part B

Time—45 minutes

Number of problems—3

No calculator is allowed for these problems.



4. A squirrel starts at building  $A$  at time  $t = 0$  and travels along a straight, horizontal wire connected to building  $B$ . For  $0 \leq t \leq 18$ , the squirrel's velocity is modeled by the piecewise-linear function defined by the graph above.
- At what times in the interval  $0 < t < 18$ , if any, does the squirrel change direction? Give a reason for your answer.
  - At what time in the interval  $0 \leq t \leq 18$  is the squirrel farthest from building  $A$ ? How far from building  $A$  is the squirrel at that time?
  - Find the total distance the squirrel travels during the time interval  $0 \leq t \leq 18$ .
  - Write expressions for the squirrel's acceleration  $a(t)$ , velocity  $v(t)$ , and distance  $x(t)$  from building  $A$  that are valid for the time interval  $7 < t < 10$ .

---

**WRITE ALL WORK IN THE EXAM BOOKLET.**

CALCULUS AB  
SECTION II, Part A  
Time—30 minutes  
Number of problems—2

A graphing calculator is required for these problems.

1. A cylindrical can of radius 10 millimeters is used to measure rainfall in Stormville. The can is initially empty, and rain enters the can during a 60-day period. The height of water in the can is modeled by the function  $S$ , where  $S(t)$  is measured in millimeters and  $t$  is measured in days for  $0 \leq t \leq 60$ . The rate at which the height of the water is rising in the can is given by  $S'(t) = 2 \sin(0.03t) + 1.5$ .
- (a) According to the model, what is the height of the water in the can at the end of the 60-day period?
- (b) According to the model, what is the average rate of change in the height of water in the can over the 60-day period? Show the computations that lead to your answer. Indicate units of measure.
- (c) Assuming no evaporation occurs, at what rate is the volume of water in the can changing at time  $t = 7$ ? Indicate units of measure.
- (d) During the same 60-day period, rain on Monsoon Mountain accumulates in a can identical to the one in Stormville. The height of the water in the can on Monsoon Mountain is modeled by the function  $M$ , where  $M(t) = \frac{1}{400}(3t^3 - 30t^2 + 330t)$ . The height  $M(t)$  is measured in millimeters, and  $t$  is measured in days for  $0 \leq t \leq 60$ . Let  $D(t) = M'(t) - S'(t)$ . Apply the Intermediate Value Theorem to the function  $D$  on the interval  $0 \leq t \leq 60$  to justify that there exists a time  $t$ ,  $0 < t < 60$ , at which the heights of water in the two cans are changing at the same rate.

2. A 12,000-liter tank of water is filled to capacity. At time  $t = 0$ , water begins to drain out of the tank at a rate modeled by  $r(t)$ , measured in liters per hour, where  $r$  is given by the piecewise-defined function

$$r(t) = \begin{cases} \frac{600t}{t+3} & \text{for } 0 \leq t \leq 5 \\ 1000e^{-0.2t} & \text{for } t > 5 \end{cases}$$

- (a) Is  $r$  continuous at  $t = 5$ ? Show the work that leads to your answer.
- (b) Find the average rate at which water is draining from the tank between time  $t = 0$  and time  $t = 8$  hours.
- (c) Find  $r'(3)$ . Using correct units, explain the meaning of that value in the context of this problem.
- (d) Write, but do not solve, an equation involving an integral to find the time  $A$  when the amount of water in the tank is 9000 liters.

WRITE ALL WORK IN THE EXAM BOOKLET.

END OF PART A OF SECTION II

© 2011 The College Board.  
Visit the College Board on the Web: [www.collegeboard.org](http://www.collegeboard.org).

GO ON TO THE NEXT PAGE.


2011 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

CALCULUS AB  
SECTION II, Part B

Time—60 minutes

Number of problems—4

No calculator is allowed for these problems.



3. The functions  $f$  and  $g$  are given by  $f(x) = \sqrt{x}$  and  $g(x) = 6 - x$ . Let  $R$  be the region bounded by the  $x$ -axis and the graphs of  $f$  and  $g$ , as shown in the figure above.
- Find the area of  $R$ .
  - The region  $R$  is the base of a solid. For each  $y$ , where  $0 \leq y \leq 2$ , the cross section of the solid taken perpendicular to the  $y$ -axis is a rectangle whose base lies in  $R$  and whose height is  $2y$ . Write, but do not evaluate, an integral expression that gives the volume of the solid.
  - There is a point  $P$  on the graph of  $f$  at which the line tangent to the graph of  $f$  is perpendicular to the graph of  $g$ . Find the coordinates of point  $P$ .

- ~~4. Consider a differentiable function  $f$  having domain all positive real numbers, and for which it is known that  $f'(x) = (4 - x)x^{-3}$  for  $x > 0$ .~~
- ~~Find the  $x$ -coordinate of the critical point of  $f$ . Determine whether the point is a relative maximum, a relative minimum, or neither for the function  $f$ . Justify your answer.~~
  - ~~Find all intervals on which the graph of  $f$  is concave down. Justify your answer.~~
  - ~~Given that  $f(1) = 2$ , determine the function  $f$ .~~

WRITE ALL WORK IN THE EXAM BOOKLET.

No calculator

2011 AP<sup>®</sup> CALCULUS AB FREE-RESPONSE QUESTIONS (Form B)

$t$ (seconds)	0	10	40	60
$B(t)$ (meters)	100	136	9	49
$v(t)$ (meters per second)	2.0	2.3	2.5	4.6

5. Ben rides a unicycle back and forth along a straight east-west track. The twice-differentiable function  $B$  models Ben's position on the track, measured in meters from the western end of the track, at time  $t$ , measured in seconds from the start of the ride. The table above gives values for  $B(t)$  and Ben's velocity,  $v(t)$ , measured in meters per second, at selected times  $t$ .
- (a) Use the data in the table to approximate Ben's acceleration at time  $t = 5$  seconds. Indicate units of measure.
- (b) Using correct units, interpret the meaning of  $\int_0^{60} |v(t)| dt$  in the context of this problem. Approximate  $\int_0^{60} |v(t)| dt$  using a left Riemann sum with the subintervals indicated by the data in the table.
- (c) For  $40 \leq t \leq 60$ , must there be a time  $t$  when Ben's velocity is 2 meters per second? Justify your answer.
- (d) A light is directly above the western end of the track. Ben rides so that at time  $t$ , the distance  $L(t)$  between Ben and the light satisfies  $(L(t))^2 = 12^2 + (B(t))^2$ . At what rate is the distance between Ben and the light changing at time  $t = 40$ ?

---

WRITE ALL WORK IN THE EXAM BOOKLET.

Name: \_\_\_\_\_

1997 AP Calculus AB:  
Section I, Part A

50 Minutes—No Calculator

Show all work!

Note: Unless otherwise specified, the domain of a function  $f$  is assumed to be the set of all real numbers  $x$  for which  $f(x)$  is a real number.

1.  $\int_1^2 (4x^3 - 6x) dx =$

- (A) 2
- (B) 4
- (C) 6
- (D) 36
- (E) 42

---

2. If  $f(x) = x\sqrt{2x-3}$ , then  $f'(x) =$

- (A)  $\frac{3x-3}{\sqrt{2x-3}}$
- (B)  $\frac{x}{\sqrt{2x-3}}$
- (C)  $\frac{1}{\sqrt{2x-3}}$
- (D)  $\frac{-x+3}{\sqrt{2x-3}}$
- (E)  $\frac{5x-6}{2\sqrt{2x-3}}$

---

3. If  $\int_a^b f(x) dx = a + 2b$ , then  $\int_a^b (f(x) + 5) dx =$

- (A)  $a + 2b + 5$       (B)  $5b - 5a$       (C)  $7b - 4a$       (D)  $7b - 5a$       (E)  $7b - 6a$

---

4. If  $f(x) = -x^3 + x + \frac{1}{x}$ , then  $f'(-1) =$

- (A) 3      (B) 1      (C) -1      (D) -3      (E) -5



1997 AP Calculus AB:  
Section I, Part A

5. The graph of  $y = 3x^4 - 16x^3 + 24x^2 + 48$  is concave down for

(A)  $x < 0$

(B)  $x > 0$

(C)  $x < -2$  or  $x > -\frac{2}{3}$

(D)  $x < \frac{2}{3}$  or  $x > 2$

(E)  $\frac{2}{3} < x < 2$

---

6.  $\frac{1}{2} \int e^{\frac{t}{2}} dt =$

(A)  $e^{-t} + C$       (B)  $e^{-\frac{t}{2}} + C$       (C)  $e^{\frac{t}{2}} + C$       (D)  $2e^{\frac{t}{2}} + C$       (E)  $e^t + C$

---

7.  $\frac{d}{dx} \cos^2(x^3) =$

(A)  $6x^2 \sin(x^3) \cos(x^3)$

(B)  $6x^2 \cos(x^3)$


(C)  $\sin^2(x^3)$

(D)  $-6x^2 \sin(x^3) \cos(x^3)$

(E)  $-2 \sin(x^3) \cos(x^3)$

1997 AP Calculus AB:  
Section I, Part A

Questions 8-9 refer to the following situation.



A bug begins to crawl up a vertical wire at time  $t = 0$ . The velocity  $v$  of the bug at time  $t$ ,  $0 \leq t \leq 8$ , is given by the function whose graph is shown above.

8. At what value of  $t$  does the bug change direction?

- (A) 2                      (B) 4                      (C) 6                      (D) 7                      (E) 8

9. What is the total distance the bug traveled from  $t = 0$  to  $t = 8$ ?

- (A) 14                      (B) 13                      (C) 11                      (D) 8                      (E) 6

10. An equation of the line tangent to the graph of  $y = \cos(2x)$  at  $x = \frac{\pi}{4}$  is

(A)  $y - 1 = -\left(x - \frac{\pi}{4}\right)$


(B)  $y - 1 = -2\left(x - \frac{\pi}{4}\right)$

(C)  $y = 2\left(x - \frac{\pi}{4}\right)$






(D)  $y = -\left(x - \frac{\pi}{4}\right)$

(E)  $y = -2\left(x - \frac{\pi}{4}\right)$

1997 AP Calculus AB:  
Section I, Part A



11. The graph of the derivative of  $f$  is shown in the figure above. Which of the following could be the graph of  $f$ ?




12. At what point on the graph of  $y = \frac{1}{2}x^2$  is the tangent line parallel to the line  $2x - 4y = 3$ ?

- (A)  $\left(\frac{1}{2}, -\frac{1}{2}\right)$     (B)  $\left(\frac{1}{2}, \frac{1}{8}\right)$     (C)  $\left(1, -\frac{1}{4}\right)$     (D)  $\left(1, \frac{1}{2}\right)$     (E)  $(2, 2)$

**1997 AP Calculus AB:  
Section I, Part A**

13. Let  $f$  be a function defined for all real numbers  $x$ . If  $f'(x) = \frac{|4-x^2|}{x-2}$ , then  $f$  is decreasing on the interval
- (A)  $(-\infty, 2)$       (B)  $(-\infty, \infty)$       (C)  $(-2, 4)$       (D)  $(-2, \infty)$       (E)  $(2, \infty)$

14. Let  $f$  be a differentiable function such that  $f(3) = 2$  and  $f'(3) = 5$ . If the tangent line to the graph of  $f$  at  $x = 3$  is used to find an approximation to a zero of  $f$ , that approximation is
- (A) 0.4      (B) 0.5      (C) 2.6      (D) 3.4      (E) 5.5



15. The graph of the function  $f$  is shown in the figure above. Which of the following statements about  $f$  is true?
- (A)  $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow b} f(x)$
- (B)  $\lim_{x \rightarrow a} f(x) = 2$
- (C)  $\lim_{x \rightarrow b} f(x) = 2$
- (D)  $\lim_{x \rightarrow b} f(x) = 1$
- (E)  $\lim_{x \rightarrow a} f(x)$  does not exist.

1997 AP Calculus AB:  
Section I, Part A

16. The area of the region enclosed by the graph of  $y = x^2 + 1$  and the line  $y = 5$  is

- (A)  $\frac{14}{3}$       (B)  $\frac{16}{3}$       (C)  $\frac{28}{3}$       (D)  $\frac{32}{3}$       (E)  $8\pi$
- 

17. If  $x^2 + y^2 = 25$ , what is the value of  $\frac{d^2y}{dx^2}$  at the point  $(4, 3)$ ?

- (A)  $-\frac{25}{27}$       (B)  $-\frac{7}{27}$       (C)  $\frac{7}{27}$       (D)  $\frac{3}{4}$       (E)  $\frac{25}{27}$
- 

18.  $\int_0^{\frac{\pi}{4}} \frac{e^{\tan x}}{\cos^2 x} dx$  is

- (A) 0      (B) 1      (C)  $e - 1$       (D)  $e$       (E)  $e + 1$
- 

19. If  $f(x) = \ln|x^2 - 1|$ , then  $f'(x) =$

- (A)  $\left| \frac{2x}{x^2 - 1} \right|$   
(B)  $\frac{2x}{|x^2 - 1|}$   
(C)  $\frac{2|x|}{x^2 - 1}$   
(D)  $\frac{2x}{x^2 - 1}$   
(E)  $\frac{1}{x^2 - 1}$

1997 AP Calculus AB:  
Section I, Part A

20. The average value of  $\cos x$  on the interval  $[-3, 5]$  is

(A)  $\frac{\sin 5 - \sin 3}{8}$

(B)  $\frac{\sin 5 - \sin 3}{2}$

(C)  $\frac{\sin 3 - \sin 5}{2}$

(D)  $\frac{\sin 3 + \sin 5}{2}$

(E)  $\frac{\sin 3 + \sin 5}{8}$

---

21.  $\lim_{x \rightarrow 1} \frac{x}{\ln x}$  is

- (A) 0                      (B)  $\frac{1}{e}$                       (C) 1                      (D)  $e$                       (E) nonexistent

---

22. What are all values of  $x$  for which the function  $f$  defined by  $f(x) = (x^2 - 3)e^{-x}$  is increasing?

- (A) There are no such values of  $x$ .  
(B)  $x < -1$  and  $x > 3$   
(C)  $-3 < x < 1$   
(D)  $-1 < x < 3$   
(E) All values of  $x$

---

23. If the region enclosed by the  $y$ -axis, the line  $y = 2$ , and the curve  $y = \sqrt{x}$  is revolved about the  $y$ -axis, the volume of the solid generated is

- (A)  $\frac{32\pi}{5}$                       (B)  $\frac{16\pi}{3}$                       (C)  $\frac{16\pi}{5}$                       (D)  $\frac{8\pi}{3}$                       (E)  $\pi$

1997 AP Calculus AB:  
Section I, Part A

24. The expression  $\frac{1}{50} \left( \sqrt{\frac{1}{50}} + \sqrt{\frac{2}{50}} + \sqrt{\frac{3}{50}} + \cdots + \sqrt{\frac{50}{50}} \right)$  is a Riemann sum approximation for

(A)  $\int_0^1 \sqrt{\frac{x}{50}} dx$

(B)  $\int_0^1 \sqrt{x} dx$

(C)  $\frac{1}{50} \int_0^1 \sqrt{\frac{x}{50}} dx$

(D)  $\frac{1}{50} \int_0^1 \sqrt{x} dx$

(E)  $\frac{1}{50} \int_0^{50} \sqrt{x} dx$

---

25.  $\int x \sin(2x) dx =$

(A)  $-\frac{x}{2} \cos(2x) + \frac{1}{4} \sin(2x) + C$

(B)  $-\frac{x}{2} \cos(2x) - \frac{1}{4} \sin(2x) + C$

(C)  $\frac{x}{2} \cos(2x) - \frac{1}{4} \sin(2x) + C$

(D)  $-2x \cos(2x) + \sin(2x) + C$

(E)  $-2x \cos(2x) - 4 \sin(2x) + C$

1997 AP Calculus AB:  
Section I, Part B

40 Minutes—~~Graphing Calculator Required~~ *permitted*

Notes: (1) The exact numerical value of the correct answer does not always appear among the choices given. When this happens, select from among the choices the number that best approximates the exact numerical value.


(2) Unless otherwise specified, the domain of a function  $f$  is assumed to be the set of all real numbers  $x$  for which  $f(x)$  is a real number.

76. If  $f(x) = \frac{e^{2x}}{2x}$ , then  $f'(x) =$

- (A) 1
- (B)  $\frac{e^{2x}(1-2x)}{2x^2}$
- (C)  $e^{2x}$
- (D)  $\frac{e^{2x}(2x+1)}{x^2}$
- (E)  $\frac{e^{2x}(2x-1)}{2x^2}$

77. The graph of the function  $y = x^3 + 6x^2 + 7x - 2 \cos x$  changes concavity at  $x =$

- (A) -1.58
- (B) -1.63
- (C) -1.67
- (D) -1.89
- (E) -2.33



78. The graph of  $f$  is shown in the figure above. If  $\int_1^3 f(x) dx = 2.3$  and  $F'(x) = f(x)$ , then  $F(3) - F(0) =$

- (A) 0.3
- (B) 1.3
- (C) 3.3
- (D) 4.3
- (E) 5.3



1997 AP Calculus AB:  
Section I, Part B

79. Let  $f$  be a function such that  $\lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h} = 5$ . Which of the following must be true?

- I.  $f$  is continuous at  $x = 2$ .
- II.  $f$  is differentiable at  $x = 2$ .
- III. The derivative of  $f$  is continuous at  $x = 2$ .

(A) I only      (B) II only      (C) I and II only      (D) I and III only      (E) II and III only

---

80. Let  $f$  be the function given by  $f(x) = 2e^{4x^2}$ . For what value of  $x$  is the slope of the line tangent to the graph of  $f$  at  $(x, f(x))$  equal to 3?

(A) 0.168      (B) 0.276      (C) 0.318      (D) 0.342      (E) 0.551

---

81. A railroad track and a road cross at right angles. An observer stands on the road 70 meters south of the crossing and watches an eastbound train traveling at 60 meters per second. At how many meters per second is the train moving away from the observer 4 seconds after it passes through the intersection?

(A) 57.60      (B) 57.88      (C) 59.20      (D) 60.00      (E) 67.40

---

82. If  $y = 2x - 8$ , what is the minimum value of the product  $xy$ ?

(A) -16      (B) -8      (C) -4      (D) 0      (E) 2

---

83. What is the area of the region in the first quadrant enclosed by the graphs of  $y = \cos x$ ,  $y = x$ , and the  $y$ -axis?

(A) 0.127      (B) 0.385      (C) 0.400      (D) 0.600      (E) 0.947

---

84. The base of a solid  $S$  is the region enclosed by the graph of  $y = \sqrt{\ln x}$ , the line  $x = e$ , and the  $x$ -axis. If the cross sections of  $S$  perpendicular to the  $x$ -axis are squares, then the volume of  $S$  is

(A)  $\frac{1}{2}$       (B)  $\frac{2}{3}$       (C) 1      (D) 2      (E)  $\frac{1}{3}(e^3 - 1)$

1997 AP Calculus AB:  
Section I, Part B

85. If the derivative of  $f$  is given by  $f'(x) = e^x - 3x^2$ , at which of the following values of  $x$  does  $f$  have a relative maximum value?

- (A)  $-0.46$       (B)  $0.20$       (C)  $0.91$       (D)  $0.95$       (E)  $3.73$
- 


86. Let  $f(x) = \sqrt{x}$ . If the rate of change of  $f$  at  $x = c$  is twice its rate of change at  $x = 1$ , then  $c =$

- (A)  $\frac{1}{4}$       (B)  $1$       (C)  $4$       (D)  $\frac{1}{\sqrt{2}}$       (E)  $\frac{1}{2\sqrt{2}}$
- 






87. At time  $t \geq 0$ , the acceleration of a particle moving on the  $x$ -axis is  $a(t) = t + \sin t$ . At  $t = 0$ , the velocity of the particle is  $-2$ . For what value  $t$  will the velocity of the particle be zero?

- (A)  $1.02$       (B)  $1.48$       (C)  $1.85$       (D)  $2.81$       (E)  $3.14$

1997 AP Calculus AB:  
Section I, Part B



88. Let  $f(x) = \int_a^x h(t) dt$ , where  $h$  has the graph shown above. Which of the following could be the graph of  $f$ ?



1997 AP Calculus AB:  
Section I, Part B

$x$	0	0.5	1.0	1.5	2.0
$f(x)$	3	3	5	8	13

89. A table of values for a continuous function  $f$  is shown above. If four equal subintervals of  $[0, 2]$  are used, which of the following is the trapezoidal approximation of  $\int_0^2 f(x) dx$ ?

- (A) 8                      (B) 12                      (C) 16                      (D) 24                      (E) 32

---

90. Which of the following are antiderivatives of  $f(x) = \sin x \cos x$ ?

I.  $F(x) = \frac{\sin^2 x}{2}$

II.  $F(x) = \frac{\cos^2 x}{2}$

III.  $F(x) = \frac{-\cos(2x)}{4}$

- (A) I only  
(B) II only  
(C) III only  
(D) I and III only  
(E) II and III only

